arXiv:hep-ph/0310002v1 30 Sep 2003

CONFINING STRINGS AND HIGH-ENERGY REACTIONS

FRANK DANIEL STEFFEN

Institute for Theoretical Physics, Bern University
Sidlerstrasse 5, CH-3012 Bern, SWITZERLAND
E-mail: steffen@itp.unibe.ch

We present the loop-loop correlation model that allows us to compute confining flux tubes and to examine their effects and manifestations in high-energy reactions.

The loop-loop correlation model (LLCM) combines perturbative gluon exchange with the non-perturbative stochastic vacuum model.[1] This combination describes the static quark-antiquark potential with color Coulomb behavior for small and confining linear rise for large source separations in agreement with lattice results. Computing the chromo-electric fields of a static color dipole in the fundamental and adjoint representation of $SU(N_c)$, confining flux tubes are obtained for large dipoles that show exact Casimir scaling.[2] Based on analytic continuation, the Euclidean LLCM provides the S-matrix element S_{DD} for high-energy dipole-dipole scattering.[2] For reactions involving large dipoles, we find clear manifestations of the confining strings. Convoluting S_{DD} with appropriate wave functions to describe dipole-hadron scattering, the confining string in the dipole can be decomposed into stringless dipoles with a given dipole number density. This allows us to calculate unintegrated gluon distributions of hadrons and photons from dipole-hadron and dipole-photon cross sections via $k_\perp$ factorization.[3]

References

1. A. I. Shoshi, F. D. Steffen, and H. J. Pirner, Nucl. Phys. A **709** (2002) 131; F. D. Steffen, arXiv:hep-ph/0301084; A. I. Shoshi, arXiv:hep-ph/0302179.
2. A. I. Shoshi, F. D. Steffen, H. G. Dosch, and H. J. Pirner, Phys. Rev. D, in press; arXiv:hep-ph/0211287.
3. A. I. Shoshi, F. D Steffen, H. G. Dosch, and H. J. Pirner, Phys. Rev. D **66** (2002) 094019; arXiv:hep-ph/0207287.